

Detect Gluon Polarisation at BNL-RHIC Through J/ψ Pair Production Process

Jiro Kodaira * and Cong-Feng Qiao†
Department of Physics, Hiroshima University
Higashi-Hiroshima 739-8526, Japan

Our recent study shows that the double spin asymmetry of J/ψ pair production is measurable in polarised $p-p$ collision at RHIC in near future. And hence it enables us to extract the polarised gluon distribution function $\Delta G(x)$.

Intrigued by the measurement of the European Muon Collaboration in polarised-target experiment [1], an enormous amount of researches have been carried out on the nucleon spin structure. The unpolarised deep inelastic scattering (DIS) experiments indicate clearly that the gluons share a large portion of the parent proton's momentum. However, what portion of proton spin is carried by gluons is still an open question. Although there are some efforts [2,3] to parameterise the polarised gluon distributions, it is still quite unclear which of these parameterisations represents the reality of nature correctly. To explore this issue, commonly believe that the study of processes other than DIS is necessary. It is now expected that the polarised proton-proton collisions at BNL relativistic heavy-ion collider(RHIC) will provide copious experimental data to unveil the polarised parton distributions.

Up to now, three main kinds of schemes in measuring the gluon polarisation are proposed, which are thought to be feasible at RHIC technically. Those are

- High- p_T Prompt Photon Production
- Jet production
- Heavy Flavor Production

There are advantages and disadvantages in each of these scenarios. For detailed discussions on this, see recent review paper of Ref. [4].

Quarkonium production and decays have long been taken as an ideal means to investigate the nature of QCD and other phenomena. Due to the approximately non-relativistic nature, the description of heavy quark and antiquark system stands as one of the simplest applications of QCD. The very clean signals of quarkonium leptonic decays enable the experimental detection with a

high precision, and therefore quarkonium plays a unique role in investigating other phenomena.

Nevertheless, theoretical description for quarkonium production is still premature, although the heavy quarkonium physics has been investigated for more than twenty years. In explaining the high- p_T J/ψ surplus production discovered by CDF group [5,6,7] at the Fermilab Tevatron, the color-octet scenario [8] was proposed based on a novel effective theory, the non-relativistic QCD(NRQCD) [9]. Having achieved the first-step success in explaining the CDF data, however, color-octet mechanism also encounters difficulties in confronting with other phenomena [10]. Due to a recent discovery [11], the extent of importance of color-octet contributions in Charmonium production remains to be unfixed.

During the past years, a series of efforts have been made on detecting polarised parton distributions through quarkonium production processes [12,13,14,15,16]. Most of these investigations are not directly applicable to RHIC physics, and they are spoiled by the uncertainties aforementioned. In recently, it is found [17] that double heavy quarkonium production in polarised proton-proton collision provides an ideal means to detect the polarised gluon distributions at RHIC, and which may at least play a supplemental role to the presently proposed program in this end. In literatures [18,19], there were discussions of detecting the gluon polarisation by means of double J/ψ production, but with main emphasis not on RHIC physics and without presenting the analytical expressions for the relative differential cross sections.

The double quarkonium production has several advantages in reducing the theoretical uncertainties mentioned above. (1) By considering double production, the relativistic corrections and color-octet uncertainties are suppressed, especially without or with a lower transverse mo-

*E-mail: kodaira@theo.phys.sci.hiroshima-u.ac.jp

†E-mail: qiao@theo.phys.sci.hiroshima-u.ac.jp




FIG. 3. Angular differential asymmetry distribution of the J/ψ pair production at RHIC with colliding energy $\sqrt{s} = 500$ GeV.

The total cross sections with different parton parameterisations are summarised in table I. The discrepancies among these results are not very large as expected.

TABLE I. Total cross sections for J/ψ pair production at RHIC with $\sqrt{s} = 500$ GeV, evaluated with different parton distributions.

	$\sigma_{\mu^+\mu^-}^{tot}$	$\sigma_{\mu^+\mu^-}(p_T > 1 \text{ GeV})$
CTEQ5L [23]	11.8 pb	7.3 pb
MRST [22]	6.5 pb	4.3 pb
GRV [24]	7.4 pb	4.7 pb

Here, the notation $\sigma_{\mu^+\mu^-}$ means that the branching ratio of $B(\psi \rightarrow \mu^+\mu^-) = 0.0588$, as the practical measuring mode to reconstruct the charmonium state, is included. From the predicted cross sections, we see that with the integrated luminosity of 800 pb^{-1} in the future run of RHIC, there will be thousands of J/ψ pair events to be detected, which can certainly give us some information on the gluon polarisation within the hadron.

In summary, it is found that the J/ψ pair production at RHIC may stand as an independent means of observing the gluon spin distributions within the hadron. The asymmetry, rather than cross sections, eliminates large amount of uncertainties which come from the non-perturbative hadronization. In the future run of RHIC, that is to say with colliding energy of 500 GeV and accumulated luminosity 800 pb^{-1} , thousands of J/ψ pair events could be detected and the gluon polarisation could be measured. With the expected upgrade of RHIC in future, the concerned process would show up to be more

important in chasing the goal of uncovering the nucleon spin structures.

The work of J.K. was supported in part by the Monbukagakaku-sho Grant-in-Aid for Scientific Research No.C-13640289. The work of C-F.Q. was supported by the Grant-in-Aid of JSPS committee.

-
- J/ψ pair production at RHIC with colliding energy $\sqrt{s} = 500$ GeV, evaluated with different parton distributions.
- [1] European Muon Collaboration, *Nucl. Phys. B* **206**, 364 (1988).
 - [2] T. Gehrmann and W.J. Stirling, *Phys. Rev. D* **53**, 6100 (1996).
 - [3] M. Glück, E. Reya and W. Vogelsang, *Phys. Lett. B* **359**, 201 (1995).
 - [4] G. Bunce, N. Saito, J. Soffer and W. Vogelsang, *Ann. Rev. Nucl. Part. Sci.* **50**, 525 (2000).
 - [5] CDF Collaboration, F. Abe *et al.*, *Phys. Rev. Lett.* **69**, 3704 (1992).
 - [6] CDF Collaboration, F. Abe *et al.*, *Phys. Rev. Lett.* **79**, 572 (1997).
 - [7] CDF Collaboration, F. Abe *et al.*, *Phys. Rev. Lett.* **79**, 578 (1997).
 - [8] E. Braaten and S. Fleming, *Phys. Rev. Lett.* **74**, 3327 (1995).
 - [9] G.T. Bodwin, E. Braaten, and G.P. Lepage, *Phys. Rev. D* **51**, 1125 (1995); **55**, 5853(E) (1997).
 - [10] For recent reviews see, for example: I. Rothstein, hep-ph/9911276; M. Krämer, hep-ph/0106120.
 - [11] Cong-Feng Qiao, hep-ph/0202227.
 - [12] A.V. Batunin and S.R. Slabospitsky, *Phys. Lett. B* **188**, 269 (1987);
 - [13] J.L. Cortes and B. Pire, *Phys. Rev. D* **38**, 3586 (1988).
 - [14] M.A. Doncheski and R.W. Robinett, *Phys. Lett. B* **248**, 188 (1990).
 - [15] T. Morii, S. Tanaka, and T. Yamanishi, *Phys. Lett. B* **372**, 165 (1996).
 - [16] R.L. Jaffe and D. Kharzeev, *Phys. Lett. B* **455**, 306 (1999).
 - [17] J. Kodaira and C.-F. Qiao, hep-ph/0207318.
 - [18] S.P. Baranov and H. Jung, *Z. Phys. C* **66**, 467 (1995).
 - [19] T. Gehrmann, *Phys. Rev. D* **53**, 5310 (1996).
 - [20] Cong-Feng Qiao, *Phys. Rev. D* **66** (2002) 057504.
 - [21] B. Humbert and P. Mery, *Z. Phys. C* **20**, 83 (1983); *Phys. Lett. B* **124**, 256 (1983).
 - [22] A.D. Martin, R.G. Roberts, W.J. Stirling and R.S Thorne, *Eur. Phys. J. C* **14**, 133 (2000).
 - [23] CTEQ Collaboration, H.L. Lai *et al.*, *Eur. Phys. J. C* **12**, 375 (2000).
 - [24] M. Glück, E. Reya and A. Vogt, *Z. Phys. C* **67**, 433 (1995).